

AD-A188 864

ACTIVITY OF ORGANOPHOSPHATE ACID ANHYDRASE IN RANGIA
CUNEATA(U) CHEMICAL RESEARCH DEVELOPMENT AND
ENGINEERING CENTER ABERDEEN PROVING GROUND MD
N A CHESTER ET AL JAN 88 CRDEC-TR-88045


1/1

UNCLASSIFIED

F/G 6/1

NL





AD-A188 864

CHEMICAL
RESEARCH,
- DEVELOPMENT &
ENGINEERING
CENTER

DTIC FILE COPY


CRDEC-TR-88045

ACTIVITY OF ORGANOPHOSPHATE ACID
ANHYDRASE IN RANGIA CUNEATA

by Nancy A. Chester
Wayne G. Landis, Ph.D.

RESEARCH DIRECTORATE

January 1988



U.S. ARMY
ARMAMENT
MUNITIONS
CHEMICAL COMMAND



Aberdeen Proving Ground Maryland 21010-5423

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

88 2 12 06 5

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) CRDEC- TR-88045			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION CRDEC		6b. OFFICE SYMBOL (if applicable) SMCCR-RST-E		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION CRDEC		8b. OFFICE SYMBOL (if applicable) SMCCR-RST-E		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO	PROJECT NO 1C463721	TASK NO DE81
			WORK UNIT ACCESSION NO		
11. TITLE (Include Security Classification) Activity of Organophosphate Acid Anhydrase in <u>Rangia Cuneata</u>					
12. PERSONAL AUTHOR(S) Chester, Nancy A., and Landis, Wayne G., Ph.D					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 86 Aug to 86 Nov		14. DATE OF REPORT (Year, Month, Day) 1988 January	
				15. PAGE COUNT 16	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Organophosphate DFP Clam		
15	06	03	Rangia cuneata DFPase		
			Mipafox Opa anhydrase		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>- Purified clam-digestive gland was used to individually test substrate solutions of DFP and Mipafox for organophosphate acid (opa) anhydrase activity. Results indicate three groups of molecular weight-estimates for substrate-specific enzymes within R. cuneata. When DFP was substrate, proteins in the 73,447 to 81,991 D and 20,157 to 43,547 D ranges were identified as opa anhydrases. With Mipafox as substrate, opa anhydrases ranging in weight from 105,026 to 138,286 D were discovered. This data suggests multiple enzymes within R. cuneata that are strictly characterized according to substrate specificity and molecular weight.</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL SANDRA J. JOHNSON			22b. TELEPHONE (Include Area Code) (301) 671-2914		22c. OFFICE SYMBOL SMCCR-SPS-T

PREFACE

The work described in this report was authorized under Project No. 1C463721DE81, Chemical Decontamination Materiel. This work was started in August 1986 and completed in November 1986.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U.S. Government purposes.

This report has been approved for release to the public.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

Blank

CONTENTS

	Page
1. INTRODUCTION	7
2. METHOD AND MATERIALS	8
2.1 Tissue Preparation	7
2.2 Column Preparation	8
2.3 Enzyme Preparation	8
3. RESULTS	8
4. DISCUSSION	9
LITERATURE CITED	11
APPENDIX. APPLE DFPASE2 PROGRAM	13

Blank

ACTIVITY OF ORGANOPHOSPHATE ACID ANHYDRASE IN RANGIA CUNEATA

1. INTRODUCTION

Enzymes capable of hydrolyzing 0,0-diisopropylfluorophosphate (DFP) and related acetylcholinesterase inhibitors, such as 0-1,2,2-trimethylpropylmethylphosphonofluoridate (soman), 3,3-dimethylbutylmethylphosphonofluoridate (Dimebu), and N,N'-diisopropylphosphorodiamidofluoridate (Mipafox), have been reported in the tissues of many animals¹⁻⁵ and have recently been renamed as organophosphate acid (opa) anhydases. Previously, two categories of DFP hydrolase enzymes, or DFPases, were partially characterized. "Squid-type" DFPase hydrolyzes DFP faster than soman,² is stable,⁶ has a molecular weight of approximately 26,000 D,⁷ is inhibited by manganous ion,⁶ experiences no inhibition of DFP hydrolysis by Mipafox,⁴ and is present in optic ganglia, giant nerve axon, hepatopancreas, and salivary gland of cephalopods.⁸ "Mazur-type" DFPase is stimulated by manganous ion, hydrolyzes soman faster than DFP,² is dimeric with a molecular weight of approximately 62,000 D,⁹ experiences inhibition of DFP hydrolysis by Mipafox,* and is unstable.² Activities resembling Mazur type are found in hog kidney, Escherichia coli, mammalian tissues, the protozoan Tetrahymena thermophila, and the clam Rangia cuneata.^{1-3,9}

However, several DFPase sources have been shown to consist of more than one type of DFPase; hence the broader term, opa anhydrase, has been brought into use (DFPase Workshop, 1987). Five isoenzymes have been recognized in T. thermophila, ranging in molecular weight from 72,000 to 96,000 Daltons, that do not fit well into the classical squid- and Mazur-type definitions.¹⁰ The use of Mipafox, which has been introduced to DFP hydrolysis studies as a tool for further enzyme identification and characterization, has provided evidence of mixtures of the DFPase types within a source. A previous study has found Mipafox to be a potent, reversible, competitive inhibitor of Mazur-type not squid-type DFPase, whether soman or DFP is used as substrate. Because of Mipafox's enzyme specificity, this study confirmed that various tissues, in particular, E. coli and squid, contain both DFPases.⁴ One recent investigation¹¹ suggested the existence of both a DFPase- and Mipafox-hydrolyzing enzyme in clam digestive gland, when enzyme-mediated hydrolysis rates appeared to be additive of rates for DFP and Mipafox. The following study continues the investigation of opa anhydrase sources with a further look at Rangia cuneata.

*Landis, W.G., Chester, N.A., Haley, M.V., Johnson, D.W., and Tauber, R.M., "Mipafox as an Inhibitor of Tetrahymena thermophila - DFPase," unpublished paper.

2. METHOD AND MATERIALS

2.1 Tissue Preparation.

Rangia cuneata (clams) were collected in sediment samples from the Chesapeake Bay near Aberdeen Proving Ground and held in ambient water at 2 °C for several hours before processing. Homogenates, 33% clam tissue by weight in Hanks balanced salt solution (HBSS) (Difco), were prepared from digestive gland pooled from 30-50 individuals and stored at 4 °C. Before testing, the tissue was diluted with Hoskin's buffer (400 mM KCl, 50 mM NaCl, and 5 mM 1,3-bis [tris hydroxymethyl methylamino] propane [bis-tris-propane] in glass distilled water, pH 7.2), and centrifuged at 1500 rpm for 20 minutes, yielding a supernatant that was placed onto a column.

2.2 Column Preparation.

Column work was conducted at 4 °C using a 2.5- x 75-cm column. Hoskin's buffer as mobile phase was passed through a sephacryl S-300 superfine grade gel matrix (Pharmacia) at a flow rate of .483 ml/min. Collection time was set at 3 minutes per fraction or 1.45 ml per fraction. The following markers (Pharmacia Gel Filtration Calibration Kit, and Bovine Albumin, Sigma) were used for column calibration: blue dextran (2,000,000 D) ferritin (440,000 D), aldolase (158,000 D), BSA (67,000 D), and ovalbumin (43,000 D). Five milliliters of clam tissue extract was placed on the column, and fractions were collected and subsequently stored at -70 °C.

2.3 Enzyme Preparation.

Hoskin's buffer was used in the assays, and all chemicals were reagent grade. Two substrate solutions were individually tested for activity, 3.0×10^{-3} M of DFP and 3.0×10^{-3} of Mipafox, by adding them to a 30 °C water-jacketed glass beaker equipped with a magnetic stirrer. Substrate hydrolysis was quantified using a fluoride electrode attached to an Orion 901 microprocessor ion analyzer that recorded fluoride concentration at 1-minute intervals. After recording spontaneous hydrolysis, enzyme-mediated hydrolysis was measured following the addition of 300 µl of partially purified tissue extract taken from sets of 10 combined fractions. Later, those fractions, in groups with the highest activities were individually tested with the same DFP and Mipafox concentrations. Determination of protein concentration of the fractions was measured using the Whitaker and Granum Ultraviolet Absorption Method.⁵ Reaction rates were calculated using the Apple-computer program, DFPase2 (Appendix).

3. RESULTS

The figure presents the activities of the fractions found to have the highest rate of hydrolysis. When fractions were assayed with DFP, proteins at three distinct molecular weight ranges were found to be active: 73,447 to 81,991 D, 20,157 to 43,547 D, and 198 to 209 D. As a result of using Mipafox as substrate, two entirely different molecular weight ranges of proteins were discovered: 105,026-128,226 D, and 2,112-3,111 D. A single fraction displayed elevated hydrolysis rates with both DFP and Mipafox as substrates. The denaturation of tissue at 70 °C for 20 minutes was found to destroy activity in the fractions whose molecular weights were below 20,000 D.

4. DISCUSSION

The digestive gland of Rangia cuneata contains substrate-specific enzymes which individually hydrolyze DFP and Mipafox. Three groups of molecular weight-estimates for heat labile enzymes allow the further characterization of clam's enzymatic activity. Previously, Rangia-DFPase was found to hydrolyze Soman greater than DFP and have a molecular weight of about 67,000-75,000 D with DFP as substrate. Exogenous Mn^{2+} stimulated soman but not DFP, hydrolysis by Clam DFPase.³ Because of these characteristics, Rangia appeared to more closely resemble the Mazur-type enzyme. However, unlike other predominantly Mazur-type tissue extracts,^{4,10} clam-digestive gland exhibits no inhibition of DFP hydrolysis by Mipafox.* Most of these extracts, taken from sources such as bacteria, protozoans, and mammals, have DFPases of molecular weight 62,000-75,000 Daltons. Our data presents a similar weight-range of 73,447-81,991 D, when DFP is substrate. This points to Mazur-type characterization; however, a second range of molecular weights, 20,157 to 43,547 D, was also determined with DFP as substrate. A weight of about 26,000 D is well within this range⁸ and is associated with the squid-type enzyme, which suggests the existence of both enzyme types.

A third set of hydrolyzing enzymes was discovered to be Mipafox-specific. Proteins ranging in weight from 105,026 to 138,286 D were discovered when the assay used Mipafox as substrate. Although a DFP hydrolyzing-enzyme of 96,000 D has been reported in T. thermophila, it may be that these high molecular weight proteins require a classification of their own. The presence of a "Mipafoxase," helps to explain a trend of increasing hydrolysis rates when the unpurified digestive gland of clams was previously assayed using separate solutions of DFP and Mipafox and a third solution containing both.* Activity rates documented in the study were lowest with Mipafox as substrate, higher with DFP, and highest with both substrates present. This trend remains unique to R. cuneata.

It has become clear that a single tissue extract, such as digestive gland from R. cuneata, may contain multiple enzymes that can no longer be categorized according to their Mazur- or squid-type characteristics. The need for a wider classification of enzymes that hydrolyze DFP and related acetylcholinesterase inhibitors has led to the renaming of these proteins as organophosphate acid anhydrases, or opa anhydrases (DFPase Workshop, 1987). In the case of the opa anhydrases of R. cuneata, each is now characterized according to substrate specificity and molecular weight.

*Chester, N.A., Anderson, P.S., and Landis, W.G., Mipafox As A Substrate for Rangia-DFPase, unpublished unclassified CPB/C report.




Figure 1. Hydrolysis of DFP and Mipafox by Clam

Opa anhydases ranging from 73,447-81,991 D and 20,157-43,547 D were identified when DFP was used as a substrate, and 105,026 to 138,286 D when Mipafox was used as a substrate. These weights were grouped according to past molecular weights: representative of Mazur-type DFPases, 62,000-75,000 D, and squid-type DFPases, about 26,000 D. Peaks at 105,296 D and 43,547 D have not been previously characterized.

LITERATURE CITED

1. Hoskin, F.C.G., Kirkish, M.A., and Steinman, K.E., "Two Enzymes for the Detoxication of Organophosphorus Compounds-Sources, Similarities, and Significance," Fund. Appl. Tox. Vol. 4, pp 5165-5172 (1984).
2. Landis, W.G., Savage, R.E., Jr., and Hoskins, F.C.G., "An Organofluorophosphate-Hydrolyzing Activity in Tetrahymena thermophila," J. Protozool. Vol. 32, Ed. 3, pp 517-519 (1985).
3. Anderson, R.S., Durst, H.D., and Landis, W.G., "DFP Hydrolysis by Tissue Extracts of the Clam, Rangia cuneata," Abstract in minutes of the Annual Meeting of the Society of Environmental Toxicology and Chemistry, November 2-5 (1986).
4. Hoskin, F.C.G., "Inhibition of a Soman- and Diisopropylphosphorofluoridate (DFP)-Detoxifying Enzyme by Mipaflox," Biochem. Pharmacol. Vol. 34, pp 2069-2072 (1985).
5. Suelter, Clarence, H., A Practical Guide to Enzymology, p 30, John Wiley and Sons, New York, NY, 1985.
6. Hoskin, F.C.G., and Long, R.J., "An Improved Sulfur Assay Applied to a Problem of Isethionate Metabolism in Squid Axon and Other Nerves," J. Neurochem. Vol. 20, pp 1317-1327 (1972).
7. Hoskin, F.C.G., and Rouch, A.H., "Hydrolysis of Nerve Gas by Squid Type Diisopropylphosphonofluoridate Hydrolyzing Enzyme on Agarose Resin," Science, Vol. 215, pp 1255-1257 (1982).
8. Hoskin, F.C.G., and Prusch, R.D., "Characterization of a DFP-Hydrolyzing Enzyme in Squid Posterior Salivary Gland by Use of Soman, DFP, and Manganous Ion," Comp. Biochem. Physiol. Vol. 75C, pp 17-20 (1983).
9. Storkebaum, W., and Witzel, H., "Study on the Enzyme Catalyzed Splitting of Triphosphates," Forschungsber. Landes Nordrhein-Westfalen. Vol. 2523, pp 1-23 (1975).
10. Landis, W.G., Durst, H.D., Savage, R.E., Jr., Haley, D.M., Haley, M.V., and Johnson, D.W., "Discovery of Multiple Organofluorophosphate Hydrolyzing Activities in The Protozoan Tetrahymena thermophila," J. Appl. Tox. Vol. 7, Ed. 1, pp 35-41 (1986).

Blank

APPLNDIX
APPLE DFPASE2 PROGRAM

Blank

Apple DEPASE2 Program

```

1  PR# 3
2  INVERSE
3  PRINT "***** *DEPASE PROGRAM 2 *****"
   *****
10 PRINT "PROGRAM FOR CALCULATION OF ENZYMATIC HYDROLYSIS RATES OF "
20 PRINT "ORGANOFLUOROPHOSPHATES BY DEPASE AS MEASURED BY FLUORIDE
   EVOLUTION"
30 PRINT "INPUTS REQUIRED ARE: TIME OF REACTION IN MINUTES, REACTION
   VOL IN ML"
40 PRINT "VOLUME OF ENZYMATIC PREP IN ML, PROTEIN CONC IN MG/ML, THE "
50 PRINT "STARTING AND FINISHING CONC OF FLUORIDE IN uM, OR THE DELTA
   IN uM"
55 PRINT " AND THE SPONTANEOUS HYDROLYSIS RATE IN uM/MIN"
60 NORMAL
61 PRINT " INPUT SPONTANEOUS HYDROLYSIS"
62 INPUT S
70 PRINT "INPUT TIME, REACTION VOLUME"
80 INPUT T,V
90 PRINT "INPUT VOL. ENZYMATIC PREP, PROTEIN CONC."
100 INPUT E,P
105 S1 = S * T
110 T1 = 60 / T
120 V1 = 1000 / V
130 E1 = 1 / E
131 PRINT "IF INPUT IS START AND FINISH CONC. KEY 0, IF DIFFERENCE KEY
   1"
132 INPUT W
133 IF W = 0 THEN GOTO 140
134 IF W = 1 THEN GOTO 135
135 PRINT "INPUT DELTA uM"
136 INPUT CT
137 GOTO 161
140 PRINT "INPUT START FLUORIDE, FINISH FLUORIDE"
150 INPUT C1, C2
160 CT = C2 - C1
161 CC = CT - S1
170 R1 = CC * (E1 * T1) / V1
171 PRINT "RUN NUMBER - 2"
172 INPUT W
174 PR# 1
175 PRINT "*****"
   *****
178 PRINT "RUN NUMBER "W" SPONTANEOUS HYDROLYSIS RATE = "S
179 PRINT "TIME = "T" VOL = "V" ENZ PREP = "E" PROTEIN = "P" uM = "CC
180 PRINT R1, "uMOLES HYDROLYZED /ML /HR"
190 R2 = R1 / 60
200 PRINT R2, "uMOLES HYDROLYZED /ML /MIN"
210 IF P = 0 THEN GOTO 240
220 P1 = 1000 / P
230 R3 = R1 * P1
240 PRINT R3, "uMOLES HYDROLYZED /uG PROTEIN /HR"

```

```

250  R4 = R3 / 60
260  PRINT R4;"uMOLES HYDROLYZED /G PROTEIN/MIN"
265  PRINT  "*****"
*****
266  PR# 0
267  PR# 3
270  PRINT "KEY 1 IF ONLY THE START AND FINISH CONC. CHANGE"
280  PRINT "KEY 2 IF TIME, VOL, PROTEIN CONC., OR VOL. ENZYME PREP
CHANGES"
285  PRINT "KEY 3 IF THE RATE OF SPONTANEOUS HYDROLYSIS ALSO CHANGES"
290  PRINT "KEY 4 IF YOU ARE FINISHED"
300  INPUT Z
310  IF Z = 1 THEN GOTO 131
320  IF Z = 2 THEN GOTO 70
330  IF Z = 3 THEN GOTO 61
335  IF Z = 4 THEN GOTO 390
390  END

```

END

FILMED

MARCH, 19 88

DTIC